



## PHYSICS STANDARD LEVEL PAPER 2

Tuesday 13 November 2012 (afternoon)

1 hour 15 minutes

0 0		C	andi	aate	sessi	on n	umb	er	
	0	0							

#### Examination code

8	8	1	2	_	6	5	0	5

## **INSTRUCTIONS TO CANDIDATES**


- Write your session number in the boxes above.
- Do not open this examination paper until instructed to do so.
- Section A: answer all questions.
- Section B: answer one question.
- Write your answers in the boxes provided.
- A calculator is required for this paper.
- A clean copy of the *Physics Data Booklet* is required for this paper.
- The maximum mark for this examination paper is [50 marks].

## **SECTION A**

Answer all questions. Write your answers in the boxes provided.


## **A1.** Data analysis question.

The movement of glaciers can be modelled by applying a load to a sample of ice.



After the load has been applied, it is observed to move downwards at a constant speed v as the ice deforms. The constant speed v is measured for different loads. The graph shows the variation of v with load W for a number of identical samples of ice.

The data points are plotted below.





(Question A1 continued)

The uncertainty in v is  $\pm 20 \,\mu\text{m s}^{-1}$  and the uncertainty in W is negligible.

(a)	(i)	On the graph opposite, draw error bars on the first and last points to show the uncertainty in $\nu$ .	[1]
	(ii)	On the graph opposite, draw the line of best-fit for the data points.	[1]
(b)	Expl	ain whether the data support the hypothesis that $v$ is directly proportional to $W$ .	[1]




(Question A1 continued)

(c) Theory suggests that the relation between v and W is

$$v=kW^3$$

where k is a constant.

To test this hypothesis a graph of  $v^{\frac{1}{3}}$  against W is plotted.



At W=5.5 N the speed is  $250\pm20 \,\mu\text{m s}^{-1}$ .

Calculate the uncertainty in  $v^{\frac{1}{3}}$  for a load of 5.5 N.


(This question continues on the following page)

[3]




(Question A1 continued)

(i)	Using the graph in (c), determine $k$ without its uncertainty.	[4]
(ii)	State an appropriate unit for your answer to (d)(i).	[1]




**A2.** This question is about magnetic fields.

A long straight vertical conductor carries an electric current. The conductor passes through a hole in a horizontal piece of paper.



(a)	State how a magnetic field arises.	[1]


(b) On the diagram below, sketch the magnetic field pattern around the long straight current-carrying conductor. The direction of the current is into the plane of the paper. [2]





(Question A2 continued)

(c) The long straight conductor is formed into a coil consisting of two separate turns, X and Y. The coil hangs with its axis vertical.



Assume that the turns of the coil each behave as a long straight conductor.

- (i) Explain why, when there is a current in the coil, the separation of X and Y decreases. [2]
- (ii) The current in the coil is 15A and the circumference of one turn is  $0.48\,\text{m}$ . In order to restore X and Y to their original separation, a mass of  $2.8\times10^{-4}\,\text{kg}$  is suspended from turn Y. Estimate the magnetic field strength at X due to Y.



Turn over

[3]

**A3.** This question is about radioactivity.

Caesium-137  $\binom{137}{55}$ Cs) is a radioactive waste product with a half-life of 30 years that is formed during the fission of uranium. Caesium-137 decays by the emission of a beta-minus  $(\beta^-)$  particle to form a nuclide of barium (Ba).

(a) State the nuclear equation for this reaction.

[2]

$$^{^{137}}_{^{55}}Cs \ \rightarrow \ ^{^{....}}_{^{....}}Ba \ + \ ^{^{0}}_{^{-1}}\beta^{-} \ + \ ......$$

(b) Determine the fraction of caesium-137 that will have decayed after 120 years.

[2]


(c) Explain, with reference to the biological effects of ionizing radiation, why it is important that humans should be shielded from the radiation emitted by caesium-137. [2]

. . . .



## **SECTION B**

This section consists of three questions: B1, B2 and B3. Answer **one** question. Write your answers in the boxes provided.

- **B1.** This question is in **two** parts. **Part 1** is about momentum change. **Part 2** is about an oscillating water column (OWC) energy converter.
  - Part 1 Momentum change

S	St	ai	te	t	h	e	la	ıv	V	0	f	c	0	n	S	e1 	ſV	⁄a _	ıt:	ic _	)T	1	0	f	1	ir	16	28	ır	r	m 	0	n	16	n	ıtı	u1	n															
																																				. <u>.</u>																	
																												•																									



(Question B1, part 1 continued)

Gravel falls vertically onto a moving horizontal conveyor belt.



			 		 	٠.		 										


(iii)	Gravel first reaches the belt at $t = 0.0$ s and continues to fall. Determine the	
	total vertical force that the gravel exerts on the conveyor belt at $t = 5.0 \mathrm{s}$ .	[3]




(Question B1, part 1 continued)

conveyor belt moves with a constant horizontal speed of 1.5 m s <sup>-1</sup> . As the gravel s on the belt, it has no horizontal speed.	
Calculate the rate of change of the kinetic energy of the gravel due to its change in horizontal speed.	,
Determine the power required to move the conveyor belt at constant speed.	
Outline why the answers to (c)(i) and (ii) are different.	
	Calculate the rate of change of the kinetic energy of the gravel due to its change in horizontal speed.  Determine the power required to move the conveyor belt at constant speed.



(Question B1 continued)


# Part 2 Oscillating water column (OWC) energy converter

(i)	State <b>one</b> other ocean-based method for power production that does not require the use of fossil fuels.
(ii)	Describe the main features of an OWC.
(iii)	Outline the energy transformations that take place in an OWC.



(Question B1, part 2 continued)

(b) (i) An OWC design has an aperture that accepts a wave width of 4.5 m. The waves at the proposed site have an average wavelength of 95 m and wave period of 8.0 s. The overall efficiency of the energy conversion of the OWC is 24%.



Assuming that the waves have a rectangular cross-section, determine the minimum wave amplitude that will be required in order for the OWC to produce a power output of  $0.10\,\mathrm{MW}$ .

Density of water=1000 kg m<sup>-3</sup>

		_							_			_					 				_			 		_				 						_		
• •	•		•	•	 •	•	 •	•	•	•	•	•	 •	•	 •	•	 •	•	•	 •	•	 •	•	 • •	•	•	•	•	 •	 •	•	•	 •	 •	•	•		
	•		•	•	 •	•	 •	•		•	•	•	 ٠	•	 •	•	 •	•	•	 •	•	 •	•	 	•		•	•	 •	 •	•	•	 •	 •	•	•		
	٠		•	•	 •	•	 •	•		•	•		 ٠	•	 •	•	 •	٠	•	 ٠	•	 •	•	 	٠		•	•	 ٠	 •	٠	•	 ٠	 •	•	•		
	•		•	•	 •		 •	٠		•	•	•	 ٠	•	 ٠		 •	٠	•	 ٠	•	 •	٠	 	•		•	•	 •	 •	•	•	 •	 •	•	•		
	•		•	•	 •	•	 •	٠		•	٠		 ٠	•	 ٠	•	 •	٠	•	 ٠	•	 •	٠	 	•		•	•	 •	 •	•	•	 •	 •	•	•		

(This question continues on the following page)




Turn over

[3]

(Question B1, part 2 continued)

(ii)	On 1	the	grid,	sketch	a	labelled	Sankey	diagram	that	represents	the	energy	
	trans	forn	nation	in this (	VC	VC.							[3]



Please do not write on this page.


Answers written on this page will not be marked.

**B2.** This question is in **two** parts. **Part 1** is about wave motion. **Part 2** is about the melting of the Pobeda ice island.

## Part 1 Wave motion

(a)	State	what	is	meant	by	the	terms	ray	and	wavefront	and	state	the	relationship	
	betwe	en the	m.												[3]


(b) The diagram shows three wavefronts, A, B and C, of a wave at a particular instant in time incident on a boundary between media X and Y. Wavefront B is also shown in medium Y.



(i) Draw a line to show wavefront C in medium Y.

[1]



(Question B2, part 1 continued)

(ii)	The refractive index of X is $n_X$ and the refractive index of Y is $n_Y$ . By making appropriate measurements, calculate $\frac{n_X}{n_Y}$ .	[3]
	wave in (b) is transverse. Describe the difference between transverse waves and gitudinal waves.	[.


(This question continues on the following page)



Turn over

(Question B2, part 1 continued)

(d) The graph below shows the variation of the velocity v with time t for one oscillating particle of medium Y.



(i)	Calculate the frequency o	f oscillation	of the particle	
(1)	Carculate the frequency of	1 OSCIII ation	of the particle.	

[2]


(ii) Identify on the graph, with the letter M, a time at which the displacement of the particle is a maximum. [1]



(Question B2, part 1 continued)

																															_
						 •											 									•	 •				
						 											 							 		-					
-																	 							 							 •
٠		 ٠															 						٠	 							 •
	•		 •		-		•	 •	•		•	٠	•	•	 •	•	 	•	 •	•	 ٠		٠	 	•	-	 •	•	 •		



(Question B2 continued)

#### Part 2 Melting of the Pobeda ice island

(a) The Pobeda ice island forms regularly when icebergs run aground near the Antarctic ice shelf. The "island", which consists of a slab of pure ice, breaks apart and melts over a period of decades. The following data are available.

> Typical dimensions of surface of island =  $70 \,\mathrm{km} \times 35 \,\mathrm{km}$ Typical height of island  $=240 \, \text{m}$ Average temperature of the island =-35°C Density of sea ice  $=920 \,\mathrm{kg} \,\mathrm{m}^{-3}$  $=3.3\times10^{5}\,\mathrm{J\,kg^{-1}}$ Specific latent heat of fusion of ice  $=2.1\times10^3\,\mathrm{J\,kg^{-1}\,K^{-1}}$ Specific heat capacity of ice

				_	gu			Wi	itl	1 1	re	efe	er	e	n	C	e 1	to	1	n	o]	le	cı	ala	ar	n	nc	oti	.01	n	ar	nd	e	ne	er	gy	7,	be	etv	W	ee	en	S	0	li	d	ic	e	aı	nd	
•	•	•	•	•		•	•	 •	•	•		•	•	•	•	•	•			•	•	•		•	•		•	•		•	•	•	•	• •	•	•	• •	•		•	•	•		•	•	•		•	•		
	•					•			•					•	•	•	•			•		•								٠			•		•	•		•										•			
		_																																																	

(ii)	Show that the energy required to melt the island to form water at 0°C is about
	$2 \times 10^{20}$ J. Assume that the top and bottom surfaces of the island are flat and that
	it has vertical sides.


(This question continues on the following page)

[3]



(Question B2, part 2 continued)

(b)


(iii)	The Sun supplies thermal energy at an average rate of $450 \mathrm{W}\mathrm{m}^{-2}$ to the surface of the island. The albedo of melting ice is 0.80. Determine an estimate of the time taken to melt the island assuming that the melted water is removed immediately	
	and that no heat is lost to the surroundings.	[3]
	gest the likely effect on the average albedo of the region in which the island was ing as a result of the melting of the Pobeda ice island.	[2]

B3. This question is in two parts. Part 1 is about a lighting system. Part 2 is about a satellite.

Part 1	Lighting	system

(a)	State Ohm's law.	[1]

(b) A lighting system is designed so that additional lamps can be added in parallel.



The diagram shows three 6 V, 9 W lamps connected in parallel to a supply of emf 6.0 V and negligible internal resistance. A fuse in the circuit melts if the current in the circuit exceeds 13 A.

(i)	Determine the maximum number of lamps that can be connected in parallel in the
	circuit without melting the fuse.


(This question continues on the following page)

[3]



(Question B3, part 1 continued)


(11)	Calculate the resistance of a lamp when operating at its normal brightness.	[1]
(iii)	By mistake, a lamp rated at 12 V, 9 W is connected in parallel with three lamps rated at 6 V, 9 W. Estimate the resistance of the circuit stating any assumption that you make.	[4]
	Estimate:	
	Assumption:	



[3]

(Question B3, part 1 continued)

(c) The diagram shows a lighting circuit which includes a light-dependent resistor (LDR).



The LDR is shielded from the lamp. Explain why the intensity of the lamp increases when darkness falls.

	 				 •					 	•				 	•			•		 	-		•		 	-				 -										-	 				
	 				 		_			 				_						_	 					 				_								_			_	 				
	 	•	 •	 •	 •	•		 •	•	 •	•	•	•	•	 	•	•	•	•	•	 •	•	•	•	•	 •	•	•	•	•	 •	•	•		•	•	•	•	 •	•	•	 	•	•		
	 	•	 •	 ٠	 •	٠	-	 •	•	 •	٠	•	•	•	 	•	•	•	٠	•	 •	•	•	•	•	 	•	•	•	•	 •	•	•	•	•	•	•	•	 •	•	-	 	٠	•		
	 	٠		 •	 -	٠		 •	•	 	•	•	•		 	•		•	-		 		•	•	-	 		•					•		•	•	•	-	 •	•	-	 	-	•		
	 									 					 				-		 					 															-	 				



(Question B3 continued)


Part 2	Satell	ite
FALL	Salen	HE

State, in words, N	Newton's universal law of gravitation.



(Question B3, part 2 continued)

(b) The diagram shows a satellite orbiting the Earth. The satellite is part of the network of global-positioning satellites (GPS) that transmit radio signals used to locate the position of receivers that are located on the Earth.



(not to scale)

When the satellite is directly overhead, the microwave signal reaches the receiver 67 ms after it leaves the satellite.

(1)	State the order of magnitude of the wavelength of microwaves.	[1]
(ii)	Calculate the height of the satellite above the surface of the Earth.	[2]



(Question B3, part 2 continued)

(c)	(i)	Explain why the satellite is accelerating towards the centre of the Earth even though its orbital speed is constant.	[2]
	(ii)	Calculate the gravitational field strength due to the Earth at the position of the satellite.	[2]
		Mass of Earth = $6.0 \times 10^{24}$ kg Radius of Earth = $6.4 \times 10^6$ m	
	(iii)	Determine the orbital speed of the satellite.	[2]



(Question B3, part 2 continued)

v )	I	ノ	:10	31	II.	Ш	10	,	П	1	П	O	u.	18	,	U	10	<i>-</i> (	וט	.U	)1(	là	,I	þ	e.	11	.0	u	C	)1	U	116	<i>-</i>	Sd	ıı	71.	111	le	•															I	. 4
														_	_	_			_	_	_	_	_	_	_	_	_			_																						_			_
			٠			•	٠	•	•	•	•	•		•			•	•										•	•	٠	•			•		٠				•	•	•			•	•		 •	٠	•	•		•		
											•							•											•	•	•			•	•		•			•		•	•		•	•		 •	•	•	•		•		
											•																			•				•						•					•	•		 •		•	•				
														. <b>.</b>													. <b>.</b>																												

