

Markscheme

November 2022

Physics

Standard level

Paper 2

11 pages

© International Baccalaureate Organization 2022

All rights reserved. No part of this product may be reproduced in any form or by any electronic or mechanical means, including information storage and retrieval systems, without the prior written permission from the IB. Additionally, the license tied with this product prohibits use of any selected files or extracts from this product. Use by third parties, including but not limited to publishers, private teachers, tutoring or study services, preparatory schools, vendors operating curriculum mapping services or teacher resource digital platforms and app developers, whether fee-covered or not, is prohibited and is a criminal offense.

More information on how to request written permission in the form of a license can be obtained from <https://ibo.org/become-an-ib-school/ib-publishing/licensing/applying-for-a-license/>.

© Organisation du Baccalauréat International 2022

Tous droits réservés. Aucune partie de ce produit ne peut être reproduite sous quelque forme ni par quelque moyen que ce soit, électronique ou mécanique, y compris des systèmes de stockage et de récupération d'informations, sans l'autorisation écrite préalable de l'IB. De plus, la licence associée à ce produit interdit toute utilisation de tout fichier ou extrait sélectionné dans ce produit. L'utilisation par des tiers, y compris, sans toutefois s'y limiter, des éditeurs, des professeurs particuliers, des services de tutorat ou d'aide aux études, des établissements de préparation à l'enseignement supérieur, des fournisseurs de services de planification des programmes d'études, des gestionnaires de plateformes pédagogiques en ligne, et des développeurs d'applications, moyennant paiement ou non, est interdite et constitue une infraction pénale.

Pour plus d'informations sur la procédure à suivre pour obtenir une autorisation écrite sous la forme d'une licence, rendez-vous à l'adresse <https://ibo.org/become-an-ib-school/ib-publishing/licensing/applying-for-a-license/>.

© Organización del Bachillerato Internacional, 2022

Todos los derechos reservados. No se podrá reproducir ninguna parte de este producto de ninguna forma ni por ningún medio electrónico o mecánico, incluidos los sistemas de almacenamiento y recuperación de información, sin la previa autorización por escrito del IB. Además, la licencia vinculada a este producto prohíbe el uso de todo archivo o fragmento seleccionado de este producto. El uso por parte de terceros —lo que incluye, a título enunciativo, editoriales, profesores particulares, servicios de apoyo académico o ayuda para el estudio, colegios preparatorios, desarrolladores de aplicaciones y entidades que presten servicios de planificación curricular u ofrezcan recursos para docentes mediante plataformas digitales—, ya sea incluido en tasas o no, está prohibido y constituye un delito.

En este enlace encontrará más información sobre cómo solicitar una autorización por escrito en forma de licencia: <https://ibo.org/become-an-ib-school/ib-publishing/licensing/applying-for-a-license/>.

Subject Details: Physics SL Paper 2 Markscheme

Mark Allocation

Candidates are required to answer ALL questions. Maximum total = [50 marks].


1. Each row in the “Question” column relates to the smallest subpart of the question.
2. The maximum mark for each question subpart is indicated in the “Total” column.
3. Each marking point in the “Answers” column is shown by means of a tick (✓) at the end of the marking point.
4. A question subpart may have more marking points than the total allows. This will be indicated by “max” written after the mark in the “Total” column.

The related rubric, if necessary, will be outlined in the “Notes” column.


5. An alternative wording is indicated in the “Answers” column by a slash (/). Either wording can be accepted.
6. An alternative answer is indicated in the “Answers” column by “OR” between the alternatives. Either answer can be accepted.
7. Words in angled brackets « » in the “Answers” column are not necessary to gain the mark.
8. **Words that are underlined are essential for the mark.**
9. **The order of marking points does not have to be as in the “Answers” column, unless stated otherwise in the “Notes” column.**
10. **If the candidate’s answer has the same “meaning” or can be clearly interpreted as being of equivalent significance, detail and validity as that in the “Answers” column then award the mark.** Where this point is considered to be particularly relevant in a question it is emphasized by **OWTTE** (or words to that effect) in the “Notes” column.
11. **Remember that many candidates are writing in a second language.** Effective communication is more important than grammatical accuracy.
12. **Occasionally, a part of a question may require an answer that is required for subsequent marking points. If an error is made in the first marking point then it should be penalized. However, if the incorrect answer is used correctly in subsequent marking points then follow through marks should be awarded.** When marking, indicate this by adding **ECF** (error carried forward) on the script. “Allow ECF” will be displayed in the “Notes” column.
13. **Do not penalize candidates for errors in units or significant figures, unless it is specifically referred to in the “Notes” column.**
14. Allow reasonable substitutions where in common usage, eg ° for rad.

Question			Answers	Notes	Total
1.	a		g OR 9.81 m s^{-2} OR acceleration of gravity/due to free fall ✓	Accept 10 m s^{-2} . Ignore sign. Do not accept bald "gravity". Accept answer that indicates tangent of the graph at time $t=0$.	1
1	b		Identification of air resistance/drag force «acting upwards» ✓ «that» increases with speed ✓ «until» weight and air resistance cancel out OR net force/acceleration becomes zero ✓	A statement as "air resistance increases with speed" scores MP1 and MP2 .	3
1	c	i	«loss in» $GPE = 3.4 \times 10^{-5} \times 9.81 \times 21 \approx 7.0 \times 10^{-3} \text{ J}$ » OR «gain in» $KE = 0.5 \times 3.4 \times 10^{-5} \times 9.0^2 \approx 1.4 \times 10^{-3} \text{ J}$ ✓ energy transferred to air $\approx 7.0 \times 10^{-3} - 1.4 \times 10^{-3} = 5.6 \times 10^{-3} \text{ J}$ ✓ any calculated answer to 2 sf ✓	Allow [1] through the use of kinematics assuming constant acceleration. Allow ECF from MP1	3
1	c	ii	«gravitational» potential energy «of the raindrop» into thermal/internal energy «of the air» ✓	Accept heat for thermal energy Accept into kinetic energy of air particles Ignore sound energy	1

Question			Answers	Notes	Total
2.	a	i	<p>energy required = $250 \times 4200 \times (30 - 15)$ ✓</p> <p>energy available = $0.30 \times 680 \times t \times A$ ✓</p> $A = \frac{250 \times 4200 \times 15}{0.30 \times 680 \times 60 \times 60} \Rightarrow 21 \text{ «m}^2\text{» OR } 22 \text{ «m}^2\text{» ✓}$	<p>Allow ECF from MP1 and MP2.</p> <p>Accept the correct use of 0.30 in either MP1 or MP2.</p>	3
2	a	ii	<p>absorbed intensity = $(1 - 0.2) \times 680 \text{ «= } 544 \text{» «W m}^{-2}\text{» OR emitted intensity} = 0.97 \times 5.67 \times 10^{-8} \times T^4$ ✓</p> $T = \sqrt[4]{\frac{544}{0.97 \times 5.67 \times 10^{-8}}} = 315 \text{ «K» ✓}$ <p>42 «°C» ✓</p>	<p>Allow ECF from MP1 and MP2.</p> <p>Allow MP1 if absorbed or emitted intensity is multiplied by area.</p>	3
2	b	i	<p>can be liquefied ✓</p> <p>has intermolecular forces / potential energy ✓</p> <p>has atoms/molecules that are not point objects / take up volume ✓</p> <p>does not follow the ideal gas law «for all T and p» ✓</p> <p>collisions between particles are non-elastic ✓</p>	<p>Accept the converse argument.</p>	1 max
2	b	ii	<p>ALTERNATIVE 1</p> <p>«constant p and V imply» $nT = \text{const}$ ✓</p> <p>T increases hence n decreases ✓</p> <p>ALTERNATIVE 2</p> <p>«constant p and n imply» V is proportional to T / air expands as it is heated ✓</p> <p>«original» air occupies a greater volume OR some air leaves through opening ✓</p>	<p>MP2 in ALT 2 must come from expansion of air, not from expansion of water.</p> <p>Award [0] for an answer based on expansion of water.</p> <p>Award [1] max for an answer based on convection currents.</p>	2
2	c		<p>solar «energy» to electrical in a photovoltaic cell AND solar to thermal in a heating panel ✓</p>		1

Question			Answers	Notes	Total
3.	a	i	<p>oscillation in antiphase ✓ smaller amplitude than P ✓</p> <p>displacement</p> 		2
3	a	ii	<p>wavelength = $\frac{2}{3} \times 0.80 = 0.53 \text{ «m»}$ ✓</p> <p>speed = $\frac{0.53}{2.8 \times 10^{-3}} = 190 \text{ «m s}^{-1}\text{»}$ ✓</p>	Allow ECF from incorrect wavelength	2

Question			Answers	Notes	Total
3	b	i	kg m s ⁻² OR m ² s ⁻² seen ✓ kg m ⁻¹ ✓	Award [2] for a BCA	2
3	b	ii	speed increases hence frequency increases ✓ by factor $\sqrt{2}$ ✓		2
3	c		travelling waves transfer energy OR standing waves don't ✓ amplitude of oscillation varies along a standing wave OR is constant along a travelling wave ✓ standing waves have nodes and antinodes OR travelling waves don't ✓ points in an internodal region have same phase in standing waves OR different phase in travelling waves ✓		2 max

Question		Answers	Notes	Total
4.	a	<p>horizontal component of any length to the left ✓ vertical component two squares long upwards ✓ E.g.</p> 	<p><i>Ignore point of application.</i> <i>Award [1] max if arrowhead not present.</i></p>	2
4	b	<p>ALTERNATIVE 1</p> <p>the net/centripetal force has constant magnitude ✓ the direction of the net/centripetal force constantly changes ✓ this is achieved by vector-adding weight and the force from the rod OR the force from the rod is vector difference of the centripetal force and weight ✓</p> <p>ALTERNATIVE 2</p> <p>at the top $F_{\text{rod}} = F_c - W$ ✓ at the bottom, $F_{\text{rod}} = F_c + W$ ✓ net F/F_c is constant so the force from the rod is different «hence is changing» ✓</p>	<p><i>Accept reference to centripetal or net force indistinctly.</i> <i>Allow reference to centripetal acceleration.</i></p>	3

Question			Answers	Notes	Total
5.	a		infinite resistance OR no current is flowing through it ✓		1
5	b	i	$\text{current} = \frac{1.47}{50.0} = 2.94 \times 10^{-2}$ «A» ✓ $r = \frac{1.49}{2.94 \times 10^{-2}} - 50.0$ OR 0.68 «Ω» ✓	<i>For MP2, allow any other correctly substituted expression for r</i>	2
5	b	ii	$1.49 \times 2.94 \times 10^{-2}$ OR $\frac{1.49^2}{50.0 + 0.68}$ OR $(2.94 \times 10^{-2})^2 (50.0 + 0.68)$ ✓ 0.0438 «W» ✓	<i>Accept use of 0.7 Ω in MP1</i>	2
5	c	i	charge/carriers are moving in a magnetic field ✓ there is a magnetic force on them / quote $F = qvB$ OR this creates a magnetic field that interacts with the external magnetic field ✓	<i>Accept electrons</i> <i>For MP2, the force must be identified as acting on charge / carriers</i>	2
5	c	ii	magnetic needle is deflected by nearby currents OR two «parallel» current-carrying wires exert a force on each other OR magnetic field due to a current can be measured directly with a probe ✓	<i>Only accept argument that refers to an observation or experiment</i>	1

Question			Answers	Notes	Total
6.	a		<p>according to $\Delta E = \Delta mc^2$ / identifies mass energy equivalence ✓</p> <p>energy is released when nucleons come together / a nucleus is formed «so nucleus has less mass than individual nucleons»</p> <p>OR</p> <p>energy is required to «completely» separate the nucleons / break apart a nucleus «so individual nucleons have more mass than nucleus» ✓</p>	<i>Accept protons and neutrons</i>	2
6	b	i	$(m_{\text{polonium}} - m_{\text{lead}} - m_{\alpha})c^2$ OR $(209.93676 - 205.92945 - 4.00151)$ OR mass difference = 5.8×10^{-3} ✓ conversion to MeV using 931.5 to give 5.4 «MeV» ✓	<i>Allow ECF from MP1 Award [2] for a BCA Award [1] for 8.6×10^{-13} J</i>	2

Question			Answers	Notes	Total
6	b	ii	<p>ALTERNATIVE 1</p> <p>energy ratio expressed in terms of momentum, e.g. $\frac{E_\alpha}{E_{\text{lead}}} = \frac{p_\alpha^2 / 2m_\alpha}{p_{\text{lead}}^2 / 2m_{\text{lead}}} \checkmark$</p> <p>$p_\alpha = p_{\text{lead}}$ hence $\frac{E_\alpha}{E_{\text{lead}}} = \frac{m_{\text{lead}}}{m_\alpha} \checkmark$</p> <p>$\frac{m_{\text{lead}}}{m_\alpha} \approx \frac{206}{4} = 51.5 \Rightarrow E_\alpha = 51.5E_{\text{lead}}$ «so α has a much greater KE»</p> <p>OR</p> <p>m_{lead} «much» greater than m_{alpha} «so α has a much greater KE» \checkmark</p> <p>ALTERNATIVE 2</p> <p>alpha particle and lead particle have equal and opposite momenta \checkmark so their velocities are inversely proportional to mass \checkmark but $\text{KE} \propto v^2$ «so α has a much greater KE» \checkmark</p>		3
6	b	iii	alpha particle \checkmark is electrically charged hence more likely to interact with electrons «in the surrounding material» \checkmark		2